arXiv:1510.06953v1 [math.GM] 19 Oct 2015

A countable set derived by fuzzy set

Huan Huang

Department of Mathematics, Jimei University, Xiamen 361021, China.

Abstract

In this paper, it shows that for each fuzzy set u on $\mathbb{R}^m$, the set $D(u)$ is at most countable. Based on this, it modifies the proof of assertion (I) in step 2 of the sufficiency part of Theorem 4.1 in paper: Characterizations of compact sets in fuzzy sets spaces with L_p metric, http://arxiv.org/abs/1509.00447.

Key words: Fuzzy sets; Level cut sets; Countable;

1 Introduction

In this paper, we show that for each fuzzy set u on $\mathbb{R}^m$, the set $D(u) = \{\alpha \in (0,1) : [u]_\alpha \nsubseteq \overline{\{u > \alpha\}}\ \}$ is at most countable.

2 Preliminaries

We introduced some basic concepts about fuzzy sets. For details, we refer to [1, 2].

We use $F(\mathbb{R}^m)$ to represent all fuzzy subsets on $\mathbb{R}^m$, i.e. functions from $\mathbb{R}^m$ to $[0,1]$. For $u \in F(\mathbb{R}^m)$, let $[u]_\alpha$ denote the α-cut of u, i.e.

$$[u]_\alpha = \begin{cases} \{x \in \mathbb{R}^m : u(x) \geq \alpha\}, & \alpha \in (0,1], \\ \operatorname{supp} u = \overline{\{x \in \mathbb{R}^m : u(x) > 0\}}, & \alpha = 0. \end{cases}$$

Let $\{u > \alpha\}$ denote the strong α-cut of u, i.e., $\{u > \alpha\} = \{x \in \mathbb{R}^m : u(x) > \alpha\}$.

Email address: hhuangjy@126.com (H. Huang).

3 Main results

Let $u \in F(\mathbb{R}^m)$, $t \in \mathbb{R}^m$ and r be a positive number in $\mathbb{R}$. Define a function $S_{u,t,r}(\cdot,\cdot) : \mathbb{S}^{m-1} \times [0,1] \to \{-\infty\} \cup \mathbb{R}$ by

$$S_{u,t,r}(e,\alpha) = \begin{cases} -\infty, & \text{if } [u]_\alpha \cap \overline{B(t,r)} = \emptyset, \\ \sup\{\langle e, x-t\rangle : x \in [u]_\alpha \cap \overline{B(t,r)}\}, & \text{if } [u]_\alpha \cap \overline{B(t,r)} \neq \emptyset, \end{cases}$$

where $\overline{B(t,r)}$ denote the closed ball $\{x \in \mathbb{R}^m : d(t,x) \leq r\}$.

We say $\alpha \in (0,1)$ is a discontinuous point of $S_{u,t,r}(e,\cdot)$ if
(i) $S_{u,t,r}(e,\alpha) \in \mathbb{R}$, and
(ii) $S_{u,t,r}(e,\beta) = -\infty$ for all $\beta > \alpha$ or $-\infty < \lim_{\beta\to\alpha+} S_{u,t,r}(e,\beta) < \lim_{\beta\to\alpha-} S_{u,t,r}(e,\beta)$.
Denote the set of all discontinuous points of $S_{u,t,r}(e,\cdot)$ by $D_{u,t,r,e}$. Then $D_{u,t,r,e}$ is at most countable because $S_{u,t,r}(e,\cdot)$ is a monotone function on $[0,1]$.

Theorem 3.1 *Let u be a fuzzy set on $\mathbb{R}^m$, t be a point in $\mathbb{R}^m$, and r be a positive real number. Then $D_{u,t,r} := \bigcup_{e\in\mathbb{S}^{m-1}} D_{u,t,r,e}$ is at most countable.*

Proof Let ϖ be a countable dense subset of $\mathbb{S}^{m-1}$. Then

$$D_{u,t,r} = \bigcup_{e\in\varpi} D_{u,t,r,e} \tag{1}$$

In fact, suppose that $\alpha \in D_{u,t,r}$, then there exists $e \in \mathbb{S}^{m-1}$ such that $\alpha \in D_{u,t,r,e}$. Hence $S(u,t,r)(e,\alpha) > -\infty$. This is equivalent to $[u]_\alpha \cap \overline{B(t,r)} \neq \emptyset$. Therefore

$$S(u,t,r)(f,\alpha) > -\infty \text{ for all } f \in \mathbb{S}^{m-1}. \tag{2}$$

To show $\alpha \in \bigcup_{e\in\varpi} D_{u,t,r,e}$, we divide the proof into two cases.

Case 1. $S(u,t,r)(e,\beta) = -\infty$ for all $\beta > \alpha$.

In this case, $[u]_\beta \cap \overline{B(t,r)} = \emptyset$ for any $\beta > \alpha$, and so $S(u,t,r)(f,\beta) = -\infty$ when $f \in \mathbb{S}^{m-1}$ and $\beta > \alpha$. Combined with (2), we know $\alpha \in D_{u,t,r,f}$ for each $f \in \mathbb{S}^{m-1}$. Thus $\alpha \in \bigcup_{e\in\varpi} D_{u,t,r,e}$.

Case 2. $-\infty < \lim_{\beta\to\alpha+} S_{u,t,r}(e,\beta) < \lim_{\beta\to\alpha-} S_{u,t,r}(e,\beta)$.

In this case, there is an $\alpha_0 > \alpha$ such that $[u]_\lambda \cap \overline{B(t,r)} \neq \emptyset$ when $\lambda \in [0,\alpha_0]$. Set

$$\xi := \lim_{\beta\to\alpha-} S_{u,t,r}(e,\beta) - \lim_{\beta\to\alpha+} S_{u,t,r}(e,\beta) > 0. \tag{3}$$

Notice that, for all $\beta \in [0,1]$ with $[u]_\beta \cap \overline{B(t,r)} \neq \emptyset$,

$|S_{u,t,r}(e,\beta) - S_{u,t,r}(f,\beta)|$

$$
\begin{aligned}
&= |\sup\{\langle e, x-t\rangle : x \in [u]_\beta \cap \overline{B(t,r)}\} - \sup\{\langle f, x-t\rangle : x \in [u]_\beta \cap \overline{B(t,r)}\}| \\
&\le \sup\{|\ \langle e-f, x-t\rangle\ |: x \in [u]_\beta \cap \overline{B(t,r)}\} \\
&\le \|e-f\| \cdot r,
\end{aligned}
$$

hence, for any $\lambda \in [0, \alpha_0]$,

$$
|\ S_{u,t,r}(e,\lambda) - S_{u,t,r}(f,\lambda)\ | \le \|e-f\| \cdot r,
$$

and so, combined with (3), we know there exists $\delta > 0$ such that, for all $f \in \mathbb{S}^{m-1} \cap B(e,\delta)$,

$$
\lim_{\beta \to \alpha-} S_{u,t,r}(f,\beta) - \lim_{\beta \to \alpha+} S_{u,t,r}(f,\beta) > \xi/2,
$$

this means that $\alpha \in D_{u,t,r,f}$ when $f \in \mathbb{S}^{m-1} \cap B(e,\delta)$. Thus there exists $g \in \varpi$ such that $\alpha \in D_{u,t,r,g}$, i.e., $\alpha \in \bigcup_{e\in\varpi} D_{u,t,r,e}$.

Now we obtain (1). Since ϖ is countable and $D_{u,t,r,e}$ is at most countable, we know $D_{u,t,r}$ is at most countable. $\square$

Remark 3.1 In the proof of Theorem 3.1, in order to show $D_{u,t,r}$ is at most countable, it proves that $D_{u,t,r} = \bigcup_{e\in\varpi} D_{u,t,r,e}$. This kind of trick was used in the proof of Lemma 4 in [3] to show a set is at most countable.

Theorem 3.2 *Let u be a fuzzy set on $\mathbb{R}^m$, then $D(u) = \{\alpha \in (0,1) : [u]_\alpha \nsubseteq \overline{\{u > \alpha\}}\ \}$ is at most countable.*

Proof If $\alpha \in D(u)$, then there is a $y \in \mathbb{R}^m$ such that $y \in [u]_\alpha$ but $y \notin \overline{\{u > \alpha\}}$. Thus,

$$
d(y, \overline{\{u > \alpha\}}) > \varepsilon > 0. \tag{4}
$$

Choose a $q \in \mathbb{Q}^m = \{(z_1, z_2, \ldots, z_m) \in \mathbb{R}^m : z_i \in \mathbb{Q},\ i = 1, 2, \ldots, m\}$ which satisfies that $\|y - q\| > 0$. We assure that $\alpha \in D_{u,q,r}$ for some $r \in Q$ with $r \ge \|y-q\|$.

In fact, let $e = (y-q)/\|y-q\|$. Then

$$
S_{u,q,r}(e,\alpha) \ge \langle e, y-q\rangle = \|y-q\|. \tag{5}
$$

If $[u]_\beta \cap \overline{B(q,r)} = \emptyset$ for any $\beta > \alpha$, then $S_{u,q,r}(e,\beta) = -\infty$ for all $\beta > \alpha$, and thus $\alpha \in D_{u,q,r,e}$.

If there exists $\beta > \alpha$ such that $[u]_\beta \cap \overline{B(q,r)} \neq \emptyset$. Pick an arbitrary $x \in \overline{\{u > \alpha\}} \cap \overline{B(q,r)}$, then, by (4),

$$
\begin{aligned}
&\|x - q\| \le r, \\
&\|x - y\| > \varepsilon.
\end{aligned}
$$

If $x = q$, then $\langle e, x - q\rangle = 0$. Suppose that $x \neq q$. Notice that

$$\frac{\langle y - q, x - q\rangle}{\|y - q\| \cdot \|x - q\|} = \cos\alpha = \frac{\|x - q\|^2 + \|y - q\|^2 - \|x - y\|^2}{2\|y - q\| \cdot \|x - q\|},$$

where α is the angle between two vectors $x - q$ and $y - q$. Thus

$$\begin{aligned}\langle e, x - q\rangle &= \frac{1}{2}\left(\frac{\|x - q\|^2}{\|y - q\|} + \|y - q\| - \frac{\|x - y\|^2}{\|y - q\|}\right)\\ &\leq \frac{1}{2}\left(\frac{r^2}{\|y - q\|} + \|y - q\| - \frac{\varepsilon^2}{\|y - q\|}\right),\end{aligned}$$

and so there exists a $\delta > 0$ such that for all $r \in [\|y - q\|, \|y - q\| + \delta)$,

$$\langle e, x - q\rangle \leq \|y - q\| - \frac{1}{4}\frac{\varepsilon^2}{\|y - q\|}. \tag{6}$$

Combined with (5) and (6), it then follows from the arbitrariness of $x \in \overline{\{u > \alpha\}} \cap \overline{B(q, r)}$ that

$$\lim_{\beta \to \alpha+} S_{u,q,r}(e, \beta) < S_{u,q,r}(e, \alpha)$$

when $r \in [\|y - q\|, \|y - q\| + \delta)$. This implies that there exists $r \in \mathbb{Q}$ such that $\alpha \in D_{u,q,r,e} \subset D_{u,q,r}$.

Now we know

$$D(u) \subseteq \bigcup_{q \in \mathbb{Q}^m, r \in \mathbb{Q}} D_{u,q,r}.$$

It then follows from Theorem 3.1 that $D(u)$ is at most countable. □

4 An application

In this section, we give an application of the main result.

The following is assertion (I) in step 2 of the sufficiency part of the proof of Theorem 4.1 in paper: Characterizations of compact sets in fuzzy sets spaces with L_p metric, see http://arxiv.org/abs/1509.00447.

(I) $P(v)$ is at most countable.

We can show this assertion in the following way.

In fact, given $\alpha \in P(v)$, it holds that $\overline{\{v > \alpha\}} \subsetneq [v]_\alpha$, and hence $\alpha \in D(v) := \{\gamma \in (0, 1) : [v]_\gamma \nsubseteq \overline{\{v > \gamma\}}\ \}$. Thus $P(v) \subseteq D(v)$. By Theorem 3.2, $D(v)$ is at most countable. So $P(v)$ is at most countable.

References

[1] C. Wu, M. Ma, The Basic of Fuzzy Analysis (in Chinese), National Defence Industry press, Beijing, 1991.

[2] P. Diamond, P. Kloeden, Metric Space of Fuzzy Sets–Theory and Application, World Scientific, Singapore, 1994.

[3] W. Trutschnig, Characterization of the sendograph-convergence of fuzzy sets by means of their L_p- and lelewise convergence, Fuzzy Sets and Systems 161 (2010) 1064-1077.